

A Level Further Mathematics A Y545 Additional Pure Mathematics Sample Question Paper

Version 2

Date – Morning/Afternoon

Time allowed: 1 hour 30 minutes

You must have:

- Printed Answer Booklet
- Formulae A Level Further Mathematics A

You may use:

- a scientific or graphical calculator



INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Complete the boxes provided on the Printed Answer Booklet with your name, centre number and candidate number.
- Answer **all** the questions.
- Write your answer to each question in the space provided in the Printed Answer Booklet.** Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION

- The total number of marks for this paper is **75**.
- The marks for each question are shown in brackets [].
- You are reminded of the need for clear presentation in your answers.**
- The Printed Answer Booklet consists of **12** pages. The Question Paper consists of **4** pages.

2

Answer **all** the questions.

- 1 A curve is given by $x = t^2 - 2 \ln t$, $y = 4t$ for $t > 0$. When the arc of the curve between the points where $t = 1$ and $t = 4$ is rotated through 2π radians about the x -axis, a surface of revolution is formed with surface area A .
Given that $A = k\pi$, where k is an integer,
- write down an integral which gives A and
 - find the value of k . [4]
- 2 Find the volume of tetrahedron OABC, where O is the origin, A = (2, 3, 1), B = (-4, 2, 5) and C = (1, 4, 4). [3]
- 3 Given $z = x \sin y + y \cos x$, show that $\frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial y^2} + z = 0$. [5]
- 4 (i) Solve the recurrence relation $u_{n+2} = 4u_{n+1} - 4u_n$ for $n \geq 0$, given that $u_0 = 1$ and $u_1 = 1$. [4]
- (ii) Show that each term of the sequence $\{u_n\}$ is an integer. [2]
- 5 **In this question you must show detailed reasoning.**
It is given that $I_n = \int_0^\pi \sin^n \theta d\theta$ for $n \geq 0$.
- (i) Prove that $I_n = \frac{n-1}{n} I_{n-2}$ for $n \geq 2$. [5]
- (ii) (a) Evaluate I_1 . [2]
- (b) Use the reduction formula to determine the exact value of $\int_0^\pi \cos^2 \theta \sin^5 \theta d\theta$. [2]
- 6 A surface S has equation $z = f(x, y)$, where $f(x, y) = 2x^2 - y^2 + 3xy + 17y$. It is given that S has a single stationary point, P .
- (i) (a) Determine the coordinates of P . [5]
- (b) Determine the nature of P . [3]
- (ii) Find the equation of the tangent plane to S at the point $Q(1, 2, 38)$. [2]

- 7 In order to rescue them from extinction, a particular species of ground-nesting birds is introduced into a nature reserve. The number of breeding pairs of these birds in the nature reserve, t years after their introduction, is an integer denoted by N_t . The initial number of breeding pairs is given by N_0 .

An initial discrete population model is proposed for N_t .

$$\text{Model I: } N_{t+1} = \frac{6}{5} N_t \left(1 - \frac{1}{900} N_t\right)$$

- (i) (a) For Model I, show that the steady state values of the number of breeding pairs are 0 and 150. [3]
- (b) Show that $N_{t+1} - N_t < 150 - N_t$ when N_t lies between 0 and 150. [3]
- (c) Hence find the long-term behaviour of the number of breeding pairs of this species of birds in the nature reserve predicted by Model I when $N_0 \in (0, 150)$. [2]

An alternative discrete population model is proposed for N_t .

$$\text{Model II: } N_{t+1} = \text{INT}\left(\frac{6}{5} N_t \left(1 - \frac{1}{900} N_t\right)\right)$$

- (ii) (a) Given that $N_0 = 8$, find the value of N_4 for each of the two models. [2]
- (b) Which of the two models gives values for N_t with the more appropriate level of precision? [1]

- 8 The set X consists of all 2×2 matrices of the form $\begin{pmatrix} x & -y \\ y & x \end{pmatrix}$, where x and y are real numbers which are not **both** zero.
- (i) (a) The matrices $\begin{pmatrix} a & -b \\ b & a \end{pmatrix}$ and $\begin{pmatrix} c & -d \\ d & c \end{pmatrix}$ are both elements of X .
 Show that $\begin{pmatrix} a & -b \\ b & a \end{pmatrix} \begin{pmatrix} c & -d \\ d & c \end{pmatrix} = \begin{pmatrix} p & -q \\ q & p \end{pmatrix}$ for some real numbers p and q to be found in terms of a, b, c and d . [2]
- (b) Prove by contradiction that p and q are not **both** zero. [5]
- (ii) Prove that X , under matrix multiplication, forms a group G .
 [You may use the result that matrix multiplication is associative.] [4]
- (iii) Determine a subgroup of G of order 17. [2]
- 9 (i) (a) Prove that $p \equiv \pm 1 \pmod{6}$ for all primes $p > 3$. [2]
- (b) Hence or otherwise prove that $p^2 - 1 \equiv 0 \pmod{24}$ for all primes $p > 3$. [3]
- (ii) Given that p is an odd prime, determine the residue of 2^{p^2-1} modulo p . [4]
- (iii) Let p and q be distinct primes greater than 3. Prove that $p^{q-1} + q^{p-1} \equiv 1 \pmod{pq}$. [5]

END OF QUESTION PAPER

Copyright Information:

OCR is committed to seeking permission to reproduce all third-party content that it uses in the assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact OCR, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.