

A Level Further Mathematics B (MEI) Y436 Further Pure with Technology Sample Question Paper

Version 2

Date – Morning/Afternoon

Time allowed: 1 hour 45 minutes

You must have:

- Printed Answer Booklet
- Formulae Further Mathematics B (MEI)

You may use:

- a scientific or graphical calculator
- Computer with appropriate software



INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Complete the boxes provided on the Printed Answer Booklet with your name, centre number and candidate number.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

COMPUTING RESOURCES

- Candidates will require access to a computer with a computer algebra system, a spreadsheet, a programming language and graph-plotting software throughout the examination.

INFORMATION

- The total mark for this paper is **60**.
- The marks for each question are shown in brackets [].
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is used. You should communicate your method with correct reasoning.
- The Printed Answer Book consists of **12** pages. The Question Paper consists of **8** pages.

2

Answer **all** the questions.

- 1** A family of curves has polar equation $r = \cos^n\left(\frac{\theta}{n}\right)$, $0 \leq \theta < n\pi$, where n is a positive *even* integer.
- (i)** (A) Sketch the curve for the cases $n = 2$ and $n = 4$. [2]
- (B) State two points which lie on every curve in the family. [1]
- (C) State one other feature common to all the curves. [1]
- (ii)** (A) Write down an integral for the length of the curve for the case $n = 4$. [2]
- (B) Evaluate the integral. [2]
- (iii)** (A) Using $t = \theta$ as the parameter, find a parametric form of the equation of the family of curves. [1]
- (B) Show that $\frac{dy}{dx} = \frac{\sin t \sin\left(\frac{t}{n}\right) - \cos t \cos\left(\frac{t}{n}\right)}{\sin t \cos\left(\frac{t}{n}\right) + \cos t \sin\left(\frac{t}{n}\right)}$. [2]
- (iv)** Hence show that there are $n + 1$ points where the tangent to the curve is parallel to the y -axis. [6]
- (v)** By referring to appropriate sketches, show that the result in part **(iv)** is true in the case $n = 4$. [2]

- 2 (i) (A) Create a program to find all the solutions to $x^2 \equiv -1 \pmod{p}$ where $0 \leq x < p$.
Write out your program in full in the Printed Answer Booklet. [5]
- (B) Use the program to find the solutions to $x^2 \equiv -1 \pmod{p}$ for the primes
- $p = 809$,
 - $p = 811$ and
 - $p = 444001$. [3]
- (ii) State Wilson's Theorem. [1]
- (iii) The following argument shows that $(4k)! \equiv ((2k)!)^2 \pmod{p}$ for the case $p = 4k + 1$.
- $$(4k)! \equiv 1 \times 2 \times 3 \times \dots \times (2k-1) \times 2k \times (2k+1) \times (2k+2) \times \dots \times (4k-1) \times 4k \pmod{p} \quad (1)$$
- $$\equiv 1 \times 2 \times 3 \times \dots \times (2k-1) \times 2k \times (-2k) \times (-(2k-1)) \times \dots \times (-2) \times (-1) \pmod{p} \quad (2)$$
- $$\equiv ((2k)!)^2 \pmod{p} \quad (3)$$
- (A) Explain why $(2k+2)$ can be written as $(-(2k-1))$ in line (2). [1]
- (B) Explain how line (3) has been obtained. [2]
- (C) Explain why, if p is a prime of the form $p = 4k + 1$, then $x^2 \equiv -1 \pmod{p}$ will have at least one solution. [1]
- (D) Hence find a solution of $x^2 \equiv -1 \pmod{29}$. [2]
- (iv) (A) Create a program that will find all the positive integers n , where $n < 1000$, such that $(n-1)! \equiv -1 \pmod{n^2}$. Write out your program in full. [3]
- (B) State the values of n obtained. [2]
- (C) A Wilson prime is a prime p such that $(p-1)! \equiv -1 \pmod{p^2}$. Write down all the Wilson primes p where $p < 1000$. [1]

- 3 This question explores the family of differential equations $\frac{dy}{dx} = \sqrt{1+ax+2y}$ for various values of the parameter a . Fig. 3 shows the tangent field in the case $a = 1$.




Fig. 3

- (i) (A) Sketch the tangent field in the case $a = -2$. [2]
- (B) Explain why the tangent field is not defined for the whole coordinate plane. [1]
- (C) Give an inequality which describes the region in which the tangent field is defined. [1]
- (D) Find a value of a such that the region for which the tangent field is defined includes the entire x -axis. [1]
- (ii) (A) For the case $a = 1$, with $y = 1$ when $x = 0$, construct a spreadsheet for the Runge-Kutta method of order 2 with formulae as follows, where $f(x, y) = \frac{dy}{dx}$.

$$k_1 = hf(x_n, y_n)$$

$$k_2 = hf(x_n + h, y_n + k_1)$$

$$y_{n+1} = y_n + \frac{1}{2}(k_1 + k_2)$$

State the formulae you have used in your spreadsheet. [3]

- (B) Use your spreadsheet to obtain the value of y correct to 4 decimal places when $x = 1$ for
- $h = 0.1$
- and
- $h = 0.05$. [2]

- (iii) (A) For the case $a=0$ find the analytical solution that passes through the point $(0, 1)$. [1]
- (B) Verify that the solution in part (iii) (A) is a solution to the differential equation. [2]
- (C) Use the solution in part (iii) (A) to find the value of y correct to 4 decimal places when $x=1$. [1]
- (iv) (A) Verify that $y = -\frac{a}{2}x + \frac{a^2}{8} - \frac{1}{2}$ is a solution for all cases when $a \leq 0$. [2]
- (B) Show that this is the only straight line solution in these cases. [4]

END OF QUESTION PAPER

BLANK PAGE

BLANK PAGE

Copyright Information:

OCR is committed to seeking permission to reproduce all third-party content that it uses in the assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact OCR, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.