

Paper 1: Pure Mathematics 1 Mark Scheme

Question	Scheme	Marks	AOs
1(a)	(i) $\frac{dy}{dx} = 12x^3 - 24x^2$	M1 A1	1.1b 1.1b
	(ii) $\frac{d^2y}{dx^2} = 36x^2 - 48x$	A1ft	1.1b
		(3)	
(b)	Substitutes $x = 2$ into their $\frac{dy}{dx} = 12 \times 2^3 - 24 \times 2^2$	M1	1.1b
	Shows $\frac{dy}{dx} = 0$ and states "hence there is a stationary point"	A1	2.1
		(2)	
(c)	Substitutes $x = 2$ into their $\frac{d^2y}{dx^2} = 36 \times 2^2 - 48 \times 2$	M1	1.1b
	$\frac{d^2y}{dx^2} = 48 > 0$ and states "hence the stationary point is a minimum"	A1ft	2.2a
		(2)	
(7 marks)			
Notes:			
(a)(i)			
M1: Differentiates to a cubic form			
A1: $\frac{dy}{dx} = 12x^3 - 24x^2$			
(a)(ii)			
A1ft: Achieves a correct $\frac{d^2y}{dx^2}$ for their $\frac{dy}{dx} = 36x^2 - 48x$			
(b)			
M1: Substitutes $x = 2$ into their $\frac{dy}{dx}$			
A1: Shows $\frac{dy}{dx} = 0$ and states "hence there is a stationary point" All aspects of the proof must be correct			
(c)			
M1: Substitutes $x = 2$ into their $\frac{d^2y}{dx^2}$			
Alternatively calculates the gradient of C either side of $x = 2$			
A1ft: For a correct calculation, a valid reason and a correct conclusion.			
Follow through on an incorrect $\frac{d^2y}{dx^2}$			

Question	Scheme	Marks	AOs
2(a)	Uses $s = r\theta \Rightarrow 3 = r \times 0.4$	M1	1.2
	$\Rightarrow OD = 7.5 \text{ cm}$	A1	1.1b
		(2)	
(b)	Uses angle $AOB = (\pi - 0.4)$ or uses radius is $(12 - '7.5')$ cm	M1	3.1a
	Uses area of sector $= \frac{1}{2}r^2\theta = \frac{1}{2} \times (12 - 7.5)^2 \times (\pi - 0.4)$	M1	1.1b
	$= 27.8\text{cm}^2$	A1ft	1.1b
		(3)	
(5 marks)			
Notes:			
(a)			
M1: Attempts to use the correct formula $s = r\theta$ with $s = 3$ and $\theta = 0.4$			
A1: $OD = 7.5 \text{ cm}$ (An answer of 7.5cm implies the use of a correct formula and scores both marks)			
(b)			
M1: $AOB = \pi - 0.4$ may be implied by the use of $AOB = \text{awrt } 2.74$ or uses radius is $(12 - \text{their '7.5'})$			
M1: Follow through on their radius $(12 - \text{their } OD)$ and their angle			
A1ft: Allow awrt 27.8 cm^2 . (Answer 27.75862562). Follow through on their $(12 - \text{their '7.5'})$ Note: Do not follow through on a radius that is negative.			

Question	Scheme	Marks	AOs
3(a)	Attempts $(x-2)^2 + (y+5)^2 = \dots$	M1	1.1b
	Centre $(2, -5)$	A1	1.1b
		(2)	
(b)	Sets $k + 2^2 + 5^2 > 0$	M1	2.2a
	$\Rightarrow k > -29$	A1ft	1.1b
		(2)	
(4 marks)			
Notes:			
(a)			
M1: Attempts to complete the square so allow $(x-2)^2 + (y+5)^2 = \dots$			
A1: States the centre is at $(2, -5)$. Also allow written separately $x = 2, y = -5$ $(2, -5)$ implies both marks			
(b)			
M1: Deduces that the right hand side of their $(x \pm \dots)^2 + (y \pm \dots)^2 = \dots$ is > 0 or ≥ 0			
A1ft: $k > -29$ Also allow $k \geq -29$ Follow through on their rhs of $(x \pm \dots)^2 + (y \pm \dots)^2 = \dots$			

Question	Scheme	Marks	AOs
4	Writes $\int \frac{t+1}{t} dt = \int 1 + \frac{1}{t} dt$ and attempts to integrate	M1	2.1
	$= t + \ln t (+c)$	M1	1.1b
	$(2a + \ln 2a) - (a + \ln a) = \ln 7$	M1	1.1b
	$a = \ln \frac{7}{2}$ with $k = \frac{7}{2}$	A1	1.1b
(4 marks)			
Notes:			
M1: Attempts to divide each term by t or alternatively multiply each term by t^{-1}			
M1: Integrates each term and knows $\int \frac{1}{t} dt = \ln t$. The $+c$ is not required for this mark			
M1: Substitutes in both limits, subtracts and sets equal to $\ln 7$			
A1: Proceeds to $a = \ln \frac{7}{2}$ and states $k = \frac{7}{2}$ or exact equivalent such as 3.5			

Question	Scheme	Marks	AOs
5	Attempts to substitute $= \frac{x+1}{2}$ into $y \Rightarrow y = 4\left(\frac{x+1}{2}\right) - 7 + \frac{6}{(x+1)}$	M1	2.1
	Attempts to write as a single fraction $y = \frac{(2x-5)(x+1)+6}{(x+1)}$	M1	2.1
	$y = \frac{2x^2-3x+1}{x+1} \quad a = -3, b = 1$	A1	1.1b

(3 marks)

Notes:

M1: Score for an attempt at substituting $t = \frac{x+1}{2}$ or equivalent into $y = 4t - 7 + \frac{3}{t}$

M1: Award this for an attempt at a single fraction with a correct common denominator.

Their $4\left(\frac{x+1}{2}\right) - 7$ term may be simplified first

A1: Correct answer only $y = \frac{2x^2-3x+1}{x+1} \quad a = -3, b = 1$

Question	Scheme	Marks	AOs
6 (a)(i)	10750 barrels	B1	3.4
(ii)	Gives a valid limitation, for example <ul style="list-style-type: none"> The model shows that the daily volume of oil extracted would become negative as t increases, which is impossible States when $t = 10, V = -1500$ which is impossible States that the model will only work for $0 \leq t \leq \frac{64}{7}$ 	B1	3.5b
		(2)	
(b)(i)	Suggests a suitable exponential model, for example $V = Ae^{kt}$	M1	3.3
	Uses $(0, 16000)$ and $(4, 9000)$ in $\Rightarrow 9000 = 16000e^{4k}$	dM1	3.1b
	$\Rightarrow k = \frac{1}{4} \ln\left(\frac{9}{16}\right)$ awrt -0.144	M1	1.1b
	$V = 16000e^{\frac{1}{4} \ln\left(\frac{9}{16}\right)t}$ or $V = 16000e^{-0.144t}$	A1	1.1b
(ii)	Uses their exponential model with $t = 3 \Rightarrow V =$ awrt 10 400 barrels	B1ft	3.4
		(5)	

(7 marks)

Notes:

(a)(i)

B1: 10750 barrels

(a)(ii)

B1: See scheme

(b)(i)

M1: Suggests a suitable exponential model, for example $V = Ae^{kt}$, $V = Ar^t$ or any other suitable function such as $V = Ae^{kt} + b$ where the candidate chooses a value for b .

dM1: Uses both $(0, 16000)$ and $(4, 9000)$ in their model.

With $V = Ae^{kt}$ candidates need to proceed to $9000 = 16000e^{4k}$

With $V = Ar^t$ candidates need to proceed to $9000 = 16000r^4$

With $V = Ae^{kt} + b$ candidates need to proceed to $9000 = (16000 - b)e^{4k} + b$ where b is given as a positive constant and $A + b = 16000$.

M1: Uses a correct method to find all constants in the model.


A1: Gives a suitable equation for the model passing through (or approximately through in the case of decimal equivalents) both values $(0, 16000)$ and $(4, 9000)$. Possible equations for the model could be for example

$$V = 16000e^{-0.144t} \quad V = 16000 \times (0.866)^t \quad V = 15800e^{-0.146t} + 200$$

(b)(ii)

B1ft: Follow through on their exponential model

Question	Scheme	Marks	AOs
7	Attempts $\vec{AC} = \vec{AB} + \vec{BC} = 2\mathbf{i} + 3\mathbf{j} + \mathbf{k} + \mathbf{i} - 9\mathbf{j} + 3\mathbf{k} = 3\mathbf{i} - 6\mathbf{j} + 4\mathbf{k}$	M1	3.1a
	Attempts to find any one length using 3-d Pythagoras	M1	2.1
	Finds all of $ AB = \sqrt{14}$, $ AC = \sqrt{61}$, $ BC = \sqrt{91}$	A1ft	1.1b
	$\cos BAC = \frac{14 + 61 - 91}{2\sqrt{14}\sqrt{61}}$	M1	2.1
	angle $BAC = 105.9^\circ$ *	A1*	1.1b
		(5)	
(5 marks)			
Notes:			
<p>M1: Attempts to find \vec{AC} by using $\vec{AC} = \vec{AB} + \vec{BC}$</p> <p>M1: Attempts to find any one length by use of Pythagoras' Theorem</p> <p>A1ft: Finds all three lengths in the triangle. Follow through on their AC</p> <p>M1: Attempts to find BAC using $\cos BAC = \frac{ AB ^2 + AC ^2 - BC ^2}{2 AB AC }$</p> <p>Allow this to be scored for other methods such as $\cos BAC = \frac{\vec{AB} \cdot \vec{AC}}{ AB AC }$</p> <p>A1*: This is a show that and all aspects must be correct. Angle $BAC = 105.9^\circ$</p>			

Question	Scheme	Marks	AOs
8 (a)	$f(3.5) = -4.8, f(4) = (+)3.1$	M1	1.1b
	Change of sign and function continuous in interval $[3.5, 4] \Rightarrow$ Root *	A1*	2.4
		(2)	
(b)	Attempts $x_1 = x_0 - \frac{f(x_0)}{f'(x_0)} \Rightarrow x_1 = 4 - \frac{3.099}{16.67}$	M1	1.1b
	$x_1 = 3.81$	A1	1.1b
	$y = \ln(2x - 5)$	(2)	
(c)		M1	3.1a
	Attempts to sketch both $y = \ln(2x - 5)$ and $y = 30 - 2x^2$		
	States that $y = \ln(2x - 5)$ meets $y = 30 - 2x^2$ in just one place, therefore $y = \ln(2x - 5) = 30 - 2x$ has just one root $\Rightarrow f(x) = 0$ has just one root	A1	2.4
		(2)	

(6 marks)

Notes:

(a)

M1: Attempts $f(x)$ at both $x = 3.5$ and $x = 4$ with at least one correct to 1 significant figure

A1*: $f(3.5)$ and $f(4)$ correct to 1 sig figure (rounded or truncated) with a correct reason and conclusion. A reason could be change of sign, or $f(3.5) \times f(4) < 0$ or similar with $f(x)$ being continuous in this interval. A conclusion could be 'Hence root' or 'Therefore root in interval'

(b)

M1: Attempts $x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$ evidenced by $x_1 = 4 - \frac{3.099}{16.67}$

A1: Correct answer only $x_1 = 3.81$

(c)

M1: For a valid attempt at showing that there is only one root. This can be achieved by

- Sketching graphs of $y = \ln(2x - 5)$ and $y = 30 - 2x^2$ on the same axes
- Showing that $f(x) = \ln(2x - 5) + 2x^2 - 30$ has no turning points
- Sketching a graph of $f(x) = \ln(2x - 5) + 2x^2 - 30$

A1: Scored for correct conclusion

Question	Scheme	Marks	AOs
9(a)	$\tan \theta + \cot \theta \equiv \frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta}$	M1	2.1
	$\equiv \frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cos \theta}$	A1	1.1b
	$\equiv \frac{1}{\frac{1}{2} \sin 2\theta}$	M1	2.1
	$\equiv 2 \operatorname{cosec} 2\theta$ *	A1*	1.1b
		(4)	
(b)	States $\tan \theta + \cot \theta = 1 \Rightarrow \sin 2\theta = 2$ AND no real solutions as $-1 \leq \sin 2\theta \leq 1$	B1	2.4
		(1)	
(5 marks)			
Notes:			
(a)			
M1: Writes $\tan \theta = \frac{\sin \theta}{\cos \theta}$ and $\cot \theta = \frac{\cos \theta}{\sin \theta}$			
A1: Achieves a correct intermediate answer of $\frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cos \theta}$			
M1: Uses the double angle formula $\sin 2\theta = 2 \sin \theta \cos \theta$			
A1*: Completes proof with no errors. This is a given answer.			
Note: There are many alternative methods. For example			
$\tan \theta + \cot \theta \equiv \tan \theta + \frac{1}{\tan \theta} \equiv \frac{\tan^2 \theta + 1}{\tan \theta} \equiv \frac{\sec^2 \theta}{\tan \theta} \equiv \frac{1}{\cos^2 \theta \times \frac{\sin \theta}{\cos \theta}} \equiv \frac{1}{\cos \theta \times \sin \theta}$ then as the			
main scheme.			
(b)			
B1: Scored for sight of $\sin 2\theta = 2$ and a reason as to why this equation has no real solutions. Possible reasons could be $-1 \leq \sin 2\theta \leq 1$and therefore $\sin 2\theta \neq 2$ or $\sin 2\theta = 2 \Rightarrow 2\theta = \arcsin 2$ which has no answers as $-1 \leq \sin 2\theta \leq 1$			

Question	Scheme	Marks	AOs
10	Use of $\frac{\sin(\theta+h) - \sin \theta}{(\theta+h) - \theta}$	B1	2.1
	Uses the compound angle identity for $\sin(A+B)$ with $A = \theta$, $B = h$ $\Rightarrow \sin(\theta+h) = \sin \theta \cos h + \cos \theta \sin h$	M1	1.1b
	Achieves $\frac{\sin(\theta+h) - \sin \theta}{h} = \frac{\sin \theta \cos h + \cos \theta \sin h - \sin \theta}{h}$	A1	1.1b
	$= \frac{\sin h}{h} \cos \theta + \left(\frac{\cos h - 1}{h} \right) \sin \theta$	M1	2.1
	Uses $h \rightarrow 0$, $\frac{\sin h}{h} \rightarrow 1$ and $\frac{\cos h - 1}{h} \rightarrow 0$ Hence the $\lim_{h \rightarrow 0} \frac{\sin(\theta+h) - \sin \theta}{(\theta+h) - \theta} = \cos \theta$ and the gradient of the chord \rightarrow gradient of the curve $\Rightarrow \frac{dy}{d\theta} = \cos \theta$ *	A1*	2.5

(5 marks)

Notes:

B1: States or implies that the gradient of the chord is $\frac{\sin(\theta+h) - \sin \theta}{h}$ or similar such as

$$\frac{\sin(\theta + \delta\theta) - \sin \theta}{\theta + \delta\theta - \theta} \text{ for a small } h \text{ or } \delta\theta$$

M1: Uses the compound angle identity for $\sin(A+B)$ with $A = \theta$, $B = h$ or $\delta\theta$

A1: Obtains $\frac{\sin \theta \cos h + \cos \theta \sin h - \sin \theta}{h}$ or equivalent

M1: Writes their expression in terms of $\frac{\sin h}{h}$ and $\frac{\cos h - 1}{h}$

A1*: Uses correct language to explain that $\frac{dy}{d\theta} = \cos \theta$

For this method they should use all of the given statements $h \rightarrow 0$, $\frac{\sin h}{h} \rightarrow 1$,

$$\frac{\cos h - 1}{h} \rightarrow 0 \text{ meaning that the } \lim_{h \rightarrow 0} \frac{\sin(\theta+h) - \sin \theta}{(\theta+h) - \theta} = \cos \theta$$

and therefore the gradient of the chord \rightarrow gradient of the curve $\Rightarrow \frac{dy}{d\theta} = \cos \theta$

Question	Scheme	Marks	AOs
10alt	Use of $\frac{\sin(\theta+h) - \sin \theta}{(\theta+h) - \theta}$	B1	2.1
	Sets $\frac{\sin(\theta+h) - \sin \theta}{(\theta+h) - \theta} = \frac{\sin\left(\theta + \frac{h}{2} + \frac{h}{2}\right) - \sin\left(\theta + \frac{h}{2} - \frac{h}{2}\right)}{h}$ and uses the compound angle identity for $\sin(A+B)$ and $\sin(A-B)$ with $A = \theta + \frac{h}{2}$, $B = \frac{h}{2}$	M1	1.1b
	Achieves $\frac{\sin(\theta+h) - \sin \theta}{h} = \frac{\left[\sin\left(\theta + \frac{h}{2}\right)\cos\left(\frac{h}{2}\right) + \cos\left(\theta + \frac{h}{2}\right)\sin\left(\frac{h}{2}\right)\right] - \left[\sin\left(\theta + \frac{h}{2}\right)\cos\left(\frac{h}{2}\right) - \cos\left(\theta + \frac{h}{2}\right)\sin\left(\frac{h}{2}\right)\right]}{h}$	A1	1.1b
	$= \frac{\sin\left(\frac{h}{2}\right)}{\frac{h}{2}} \times \cos\left(\theta + \frac{h}{2}\right)$	M1	2.1
	Uses $h \rightarrow 0$, $\frac{h}{2} \rightarrow 0$ hence $\frac{\sin\left(\frac{h}{2}\right)}{\frac{h}{2}} \rightarrow 1$ and $\cos\left(\theta + \frac{h}{2}\right) \rightarrow \cos \theta$ Therefore the $\lim_{h \rightarrow 0} \frac{\sin(\theta+h) - \sin \theta}{(\theta+h) - \theta} = \cos \theta$ and the gradient of the chord \rightarrow gradient of the curve $\Rightarrow \frac{dy}{d\theta} = \cos \theta$ *	A1*	2.5

(5 marks)

Additional notes:

A1*: Uses correct language to explain that $\frac{dy}{d\theta} = \cos \theta$. For this method they should use the

(adapted) given statement $h \rightarrow 0$, $\frac{h}{2} \rightarrow 0$ hence $\frac{\sin\left(\frac{h}{2}\right)}{\frac{h}{2}} \rightarrow 1$ with $\cos\left(\theta + \frac{h}{2}\right) \rightarrow \cos \theta$

meaning that the $\lim_{h \rightarrow 0} \frac{\sin(\theta+h) - \sin \theta}{(\theta+h) - \theta} = \cos \theta$ and therefore the gradient of the

chord \rightarrow gradient of the curve $\Rightarrow \frac{dy}{d\theta} = \cos \theta$

Question	Scheme	Marks	AOs
11(a)	Sets $H = 0 \Rightarrow 1.8 + 0.4d - 0.002d^2 = 0$	M1	3.4
	Solves using an appropriate method, for example $d = \frac{-0.4 \pm \sqrt{(0.4)^2 - 4(-0.002)(1.8)}}{2 \times -0.002}$	dM1	1.1b
	Distance = awrt 204(m) only	A1	2.2a
		(3)	
(b)	States the initial height of the arrow above the ground.	B1	3.4
		(1)	
(c)	$1.8 + 0.4d - 0.002d^2 = -0.002(d^2 - 200d) + 1.8$	M1	1.1b
	$= -0.002((d - 100)^2 - 10000) + 1.8$	M1	1.1b
	$= 21.8 - 0.002(d - 100)^2$	A1	1.1b
		(3)	
(d)	(i) 22.1 metres	B1ft	3.4
	(ii) 100 metres	B1ft	3.4
		(2)	
			(9 marks)
Notes:			
(a)			
M1: Sets $H = 0 \Rightarrow 1.8 + 0.4d - 0.002d^2 = 0$			
M1: Solves using formula, which if stated must be correct, by completing square (look for $(d - 100)^2 = 10900 \Rightarrow d = ..$) or even allow answers coming from a graphical calculator			
A1: Awrt 204 m only			
(b)			
B1: States it is the initial height of the arrow above the ground. Do not allow "it is the height of the archer"			
(c)			
M1: Score for taking out a common factor of -0.002 from at least the d^2 and d terms			
M1: For completing the square for their $(d^2 - 200d)$ term			
A1: $= 21.8 - 0.002(d - 100)^2$ or exact equivalent			
(d)			
B1ft: For their '21.8+0.3' =22.1m			
B1ft: For their 100m			

Question	Scheme	Marks	AOs
12 (a)	$N = aT^b \Rightarrow \log_{10} N = \log_{10} a + \log_{10} T^b$	M1	2.1
	$\Rightarrow \log_{10} N = \log_{10} a + b \log_{10} T$ so $m = b$ and $c = \log_{10} a$	A1	1.1b
		(2)	
(b)	Uses the graph to find either a or b $a = 10^{\text{intercept}}$ or $b = \text{gradient}$	M1	3.1b
	Uses the graph to find both a and b $a = 10^{\text{intercept}}$ and $b = \text{gradient}$	M1	1.1b
	Uses $T = 3$ in $N = aT^b$ with their a and b	M1	3.1b
	Number of microbes ≈ 800	A1	1.1b
		(4)	
(c)	$N = 1000000 \Rightarrow \log_{10} N = 6$	M1	3.4
	We cannot 'extrapolate' the graph and assume that the model still holds	A1	3.5b
		(2)	
(d)	States that ' a ' is the number of microbes 1 day after the start of the experiment	B1	3.2a
		(1)	
			(9 marks)

Question 12 continued**Notes:****(a)****M1:** Takes logs of both sides and shows the addition law**M1:** Uses the power law, writes $\log_{10} N = \log_{10} a + b \log_{10} T$ and states $m = b$ **and** $c = \log_{10} a$ **(b)****M1:** Uses the graph to find either a or b $a = 10^{\text{intercept}}$ **or** $b = \text{gradient}$. This would be implied by the sight of $b = 2.3$ **or** $a = 10^{1.8} \approx 63$ **M1:** Uses the graph to find both a and b $a = 10^{\text{intercept}}$ **and** $b = \text{gradient}$. This would be implied by the sight of $b = 2.3$ **and** $a = 10^{1.8} \approx 63$ **M1:** Uses $T = 3 \Rightarrow N = aT^b$ with their a and b . This is implied by an attempt at $63 \times 3^{2.3}$ **A1:** Accept a number of microbes that are approximately 800. Allow 800 ± 150 following correct work.

There is an alternative to this using a graphical approach.

M1: Finds the value of $\log_{10} T$ from $T = 3$. Accept as $T = 3 \Rightarrow \log_{10} T \approx 0.48$ **M1:** Then using the line of best fit finds the value of $\log_{10} N$ from their "0.48"Accept $\log_{10} N \approx 2.9$ **M1:** Finds the value of N from their value of $\log_{10} N$ $\log_{10} N \approx 2.9 \Rightarrow N = 10^{2.9}$ **A1:** Accept a number of microbes that are approximately 800. Allow 800 ± 150 following correct work**(c)****M1** For using $N = 1000000$ and stating that $\log_{10} N = 6$ **A1:** Statement to the effect that "we only have information for values of $\log N$ between 1.8 and 4.5 so we cannot be certain that the relationship still holds". "We cannot extrapolate with any certainty, we could only interpolate"

There is an alternative approach that uses the formula.

M1: Use $N = 1000000$ in their $N = 63 \times T^{2.3} \Rightarrow \log_{10} T = \frac{\log_{10} \left(\frac{1000000}{63} \right)}{2.3} \approx 1.83$.**A1:** The reason would be similar to the main scheme as we only have $\log_{10} T$ values from 0 to 1.2. We cannot 'extrapolate' the graph and assume that the model still holds**(d)****B1:** Allow a numerical explanation $T = 1 \Rightarrow N = a1^b \Rightarrow N = a$ giving a is the value of N at $T = 1$

Question	Scheme	Marks	AOs
13(a)	Attempts $\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$	M1	1.1b
	$\frac{dy}{dx} = \frac{\sqrt{3} \sin 2t}{\sin t} \quad (= 2\sqrt{3} \cos t)$	A1	1.1b
		(2)	
(b)	Substitutes $t = \frac{2\pi}{3}$ in $\frac{dy}{dx} = \frac{\sqrt{3} \sin 2t}{\sin t} = (-\sqrt{3})$	M1	2.1
	Uses gradient of normal = $-\frac{1}{dy/dx} = \left(\frac{1}{\sqrt{3}}\right)$	M1	2.1
	Coordinates of $P = \left(-1, -\frac{\sqrt{3}}{2}\right)$	B1	1.1b
	Correct form of normal $y + \frac{\sqrt{3}}{2} = \frac{1}{\sqrt{3}}(x+1)$	M1	2.1
	Completes proof $\Rightarrow 2x - 2\sqrt{3}y - 1 = 0$ *	A1*	1.1b
		(5)	
(c)	Substitutes $x = 2\cos t$ and $y = \sqrt{3} \cos 2t$ into $2x - 2\sqrt{3}y - 1 = 0$	M1	3.1a
	Uses the identity $\cos 2t = 2\cos^2 t - 1$ to produce a quadratic in $\cos t$	M1	3.1a
	$\Rightarrow 12\cos^2 t - 4\cos t - 5 = 0$	A1	1.1b
	Finds $\cos t = \frac{5}{6}, \frac{1}{2}$	M1	2.4
	Substitutes their $\cos t = \frac{5}{6}$ into $x = 2\cos t, y = \sqrt{3} \cos 2t,$	M1	1.1b
	$Q = \left(\frac{5}{3}, \frac{7}{18}\sqrt{3}\right)$	A1	1.1b
		(6)	
(13 marks)			

Question 13 continued**Notes:****(a)**

M1: Attempts $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$ and achieves a form $k \frac{\sin 2t}{\sin t}$ Alternatively candidates may apply the

double angle identity for $\cos 2t$ and achieve a form $k \frac{\sin t \cos t}{\sin t}$

A1: Scored for a correct answer, either $\frac{\sqrt{3} \sin 2t}{\sin t}$ or $2\sqrt{3} \cos t$

(b)

M1: For substituting $t = \frac{2\pi}{3}$ in their $\frac{dy}{dx}$ which must be in terms of t

M1: Uses the gradient of the normal is the negative reciprocal of the value of $\frac{dy}{dx}$. This may be seen in the equation of l .

B1: States or uses (in their tangent or normal) that $P = \left(-1, -\frac{\sqrt{3}}{2}\right)$

M1: Uses their numerical value of $-1/\frac{dy}{dx}$ with their $\left(-1, -\frac{\sqrt{3}}{2}\right)$ to form an equation of the normal at P

A1*: This is a proof and all aspects need to be correct. Correct answer only $2x - 2\sqrt{3}y - 1 = 0$

(c)

M1: For substituting $x = 2\cos t$ and $y = \sqrt{3} \cos 2t$ into $2x - 2\sqrt{3}y - 1 = 0$ to produce an equation in t . Alternatively candidates could use $\cos 2t = 2\cos^2 t - 1$ to set up an equation of the form $y = Ax^2 + B$.

M1: Uses the identity $\cos 2t = 2\cos^2 t - 1$ to produce a quadratic equation in $\cos t$
In the alternative method it is for combining their $y = Ax^2 + B$ with $2x - 2\sqrt{3}y - 1 = 0$ to get an equation in just one variable

A1: For the correct quadratic equation $12\cos^2 t - 4\cos t - 5 = 0$

Alternatively the equations in x and y are $3x^2 - 2x - 5 = 0$ $12\sqrt{3}y^2 + 4y - 7\sqrt{3} = 0$

M1: Solves the quadratic equation in $\cos t$ (or x or y) and rejects the value corresponding to P .

M1: Substitutes their $\cos t = \frac{5}{6}$ or their $t = \arccos\left(\frac{5}{6}\right)$ in $x = 2\cos t$ and $y = \sqrt{3} \cos 2t$

If a value of x or y has been found it is for finding the other coordinate.

A1: $Q = \left(\frac{5}{3}, \frac{7}{18}\sqrt{3}\right)$. Allow $x = \frac{5}{3}, y = \frac{7}{18}\sqrt{3}$ but do not allow decimal equivalents.

Question	Scheme	Marks	AOs
14(a)	Uses or implies $h = 0.5$	B1	1.1b
	For correct form of the trapezium rule =	M1	1.1b
	$\frac{0.5}{2} \{3 + 2.2958 + 2(2.3041 + 1.9242 + 1.9089)\} = 4.393$	A1	1.1b
		(3)	
(b)	Any valid statement reason, for example <ul style="list-style-type: none"> • Increase the number of strips • Decrease the width of the strips • Use more trapezia 	B1	2.4
		(1)	
(c)	For integration by parts on $\int x^2 \ln x \, dx$	M1	2.1
	$= \frac{x^3}{3} \ln x - \int \frac{x^2}{3} \, dx$	A1	1.1b
	$\int -2x + 5 \, dx = -x^2 + 5x \quad (+c)$	B1	1.1b
	All integration attempted and limits used		
	Area of $S = \int_1^3 \frac{x^2 \ln x}{3} - 2x + 5 \, dx = \left[\frac{x^3}{9} \ln x - \frac{x^3}{27} - x^2 + 5x \right]_{x=1}^{x=3}$	M1	2.1
	Uses correct ln laws, simplifies and writes in required form	M1	2.1
	Area of $S = \frac{28}{27} + \ln 27 \quad (a = 28, b = 27, c = 27)$	A1	1.1b
	(6)		
(10 marks)			

Question 14 continued**Notes:****(a)**

B1: States or uses the strip width $h = 0.5$. This can be implied by the sight of $\frac{0.5}{2}\{\dots\}$ in the trapezium rule

M1: For the correct form of the bracket in the trapezium rule. Must be y values rather than x values $\{\text{first } y \text{ value} + \text{last } y \text{ value} + 2 \times (\text{sum of other } y \text{ values})\}$

A1: 4.393

(b)

B1: See scheme

(c)

M1: Uses integration by parts the right way around.

Look for $\int x^2 \ln x \, dx = Ax^3 \ln x - \int Bx^2 \, dx$

A1: $\frac{x^3}{3} \ln x - \int \frac{x^2}{3} \, dx$

B1: Integrates the $-2x + 5$ term correctly $= -x^2 + 5x$

M1: All integration completed and limits used

M1: Simplifies using \ln law(s) to a form $\frac{a}{b} + \ln c$

A1: Correct answer only $\frac{28}{27} + \ln 27$

Question	Scheme	Marks	AOs
15(a)	Attempts to differentiate using the quotient rule or otherwise	M1	2.1
	$f'(x) = \frac{e^{\sqrt{2}x-1} \times 8 \cos 2x - 4 \sin 2x \times \sqrt{2} e^{\sqrt{2}x-1}}{(e^{\sqrt{2}x-1})^2}$	A1	1.1b
	Sets $f'(x) = 0$ and divides/ factorises out the $e^{\sqrt{2}x-1}$ terms	M1	2.1
	Proceeds via $\frac{\sin 2x}{\cos 2x} = \frac{8}{4\sqrt{2}}$ to $\Rightarrow \tan 2x = \sqrt{2}$ *	A1*	1.1b
		(4)	
(b)	(i) Solves $\tan 4x = \sqrt{2}$ and attempts to find the 2 nd solution	M1	3.1a
	$x = 1.02$	A1	1.1b
	(ii) Solves $\tan 2x = \sqrt{2}$ and attempts to find the 1 st solution	M1	3.1a
	$x = 0.478$	A1	1.1b
		(4)	
(8 marks)			

Notes:

(a)

M1: Attempts to differentiate by using the quotient rule with $u = 4 \sin 2x$ and $v = e^{\sqrt{2}x-1}$ or alternatively uses the product rule with $u = 4 \sin 2x$ and $v = e^{1-\sqrt{2}x}$

A1: For achieving a correct $f'(x)$. For the product rule

$$f'(x) = e^{1-\sqrt{2}x} \times 8 \cos 2x + 4 \sin 2x \times -\sqrt{2} e^{1-\sqrt{2}x}$$

M1: This is scored for cancelling/ factorising out the exponential term. Look for an equation in just $\cos 2x$ and $\sin 2x$

A1*: Proceeds to $\tan 2x = \sqrt{2}$. This is a given answer.

(b) (i)

M1: Solves $\tan 4x = \sqrt{2}$ attempts to find the 2nd solution. Look for $x = \frac{\pi + \arctan \sqrt{2}}{4}$

Alternatively finds the 2nd solution of $\tan 2x = \sqrt{2}$ and attempts to divide by 2

A1: Allow awrt $x = 1.02$. The correct answer, with no incorrect working scores both marks

(b)(ii)

M1: Solves $\tan 2x = \sqrt{2}$ attempts to find the 1st solution. Look for $x = \frac{\arctan \sqrt{2}}{2}$

A1: Allow awrt $x = 0.478$. The correct answer, with no incorrect working scores both marks